

Review

## Review of ICT-based services for identified unmet needs in people with dementia

S. Lauriks<sup>a,\*</sup>, A. Reinersmann<sup>a</sup>, H.G. Van der Roest<sup>a</sup>, F.J.M. Meiland<sup>a</sup>,  
R.J. Davies<sup>b</sup>, F. Moelaert<sup>c</sup>, M.D. Mulvenna<sup>b</sup>, C.D. Nugent<sup>b</sup>, R.M. Dröes<sup>d,\*</sup>

<sup>a</sup>Regional Mental Health Institute GGZ-Buitendamstel Geestgronden/Alzheimer Center, VU University Medical Center, Amsterdam, The Netherlands

<sup>b</sup>School of Computing and Mathematics, University of Ulster, United Kingdom

<sup>c</sup>Telematica Institute, The Netherlands

<sup>d</sup>Department of Psychiatry/Alzheimer Center, VU University Medical Center, Amsterdam, The Netherlands

Received 27 March 2007; received in revised form 19 July 2007; accepted 23 July 2007

### Abstract

Some of the needs that people with dementia and their informal carers currently perceive as insufficiently met by regular care and support services might be alleviated, or even be met, using modern Information and Communication Technology (ICT). The study described in this paper was designed to provide an insight into the state of the art in ICT solutions that could contribute to meet the most frequently mentioned unmet needs by people with dementia and their informal carers. These needs can be summarized as (1) the need for general and personalized information; (2) the need for support with regard to symptoms of dementia; (3) the need for social contact and company; and (4) the need for health monitoring and perceived safety. Databases that were searched include: PubMed, Cinahl, Psychinfo, Google (Scholar), INSPEC and IEEE. In total 22 websites and 46 publications were included that satisfied the following criteria: the article reports on people with dementia and/or their informal carers and discusses an ICT-device that has been tested within the target group and has proven to be helpful. Within the first need area 18 relevant websites and three studies were included; within the second need area 4 websites and 20 publications were included. Within the third and fourth need area 11 and 12 publications were included respectively. Most articles reported on uncontrolled studies. It is concluded that the informational websites offer helpful information for carers but seem less attuned to the person with dementia and do not offer personalized information. ICT solutions aimed at compensating for disabilities, such as memory problems and daily activities demonstrate that people with mild to moderate dementia are capable of handling simple electronic equipment and can benefit from it in terms of more confidence and enhanced positive effect. Instrumental ICT-support for coping with behavioral and psychological changes in dementia is relatively disregarded as yet, while support for social contact can be effectively realized through, for example, simplified (mobile) phones or videophones or (entertainment) robots. GPS technology and monitoring systems are proven to result in enhanced feelings of safety and less fear and anxiety. Though these results are promising, more controlled studies in which the developed ICT solutions are tested in real life situations are needed before implementing them in the care for people with dementia. It is recommended that future studies also focus on the integration of the current techniques and solutions.

© 2007 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Dementia; Unmet needs; Technology; Support

\* Corresponding authors at: Department of Psychiatry, VU University Medical Center, Valeriusplein 9, 1075 BG Amsterdam, The Netherlands. Tel.: +31 20 7885454.

E-mail addresses: [s.lauriks@vumc.nl](mailto:s.lauriks@vumc.nl) (S. Lauriks), [rm.droes@vumc.nl](mailto:rm.droes@vumc.nl) (R.M. Dröes).

## 1. Introduction

As the population in our society is ageing, the number of people with chronic diseases is growing. To serve this growing group in an effective and efficient way, more specific knowledge of the needs of people with chronic diseases, as well as of individual variations in needs within different groups of people with chronic disease, is required. One of these groups is people with dementia, whose numbers are expected to double over the next two decades (Ferri et al., 2005). Most of the people with dementia live in the community and are supported by family and friends and professional carers and services. A minority live in institutions such as homes for the elderly, sheltered housing projects and nursing homes (Health Council of the Netherlands, 2002; Ferri et al., 2005).

People with dementia have many needs during the progression of their disease, varying from memory support in mild dementia to support in almost all aspects of daily functioning in severe dementia. Family carers, neighbors and friends meet some of these needs, while professional carers meet others. However, despite the efforts of these informal and formal carers, not all the needs of people with dementia can be met. The reasons for this include the limited time that both informal and formal carers have available, and the lack of, or limited availability of, professional services attuned specifically to the concrete needs of individual people with dementia.

A recent gap analysis in two regions of the Netherlands (Amsterdam and Nijmegen) showed that various types of services are available for all of the most frequently mentioned unmet needs (Dröes et al., 2005b). Several possible reasons were mentioned to explain the fact that these needs are still perceived as unmet by people with dementia and their carers. For example, patient and carer (or even professional referrers) are not aware of the service offering that is available; they experience a threshold for using professional services; there are mismatches between the way in which services are offered and the individual needs of patients and/or carers. These reasons are confirmed in the literature, where Coe and Neufeld (1999) conclude that most people do not know where to go for help and that they have to take the initiative to find it. As one carer expresses: “There is help if you just go after it. The help doesn’t find you. You have to go after the help.” (p. 575). Cox (1997) and Brodaty et al. (2005) reported that carers who did not use respite services felt they did not really need them, while Koffman and Taylor (1997) observed that carers experienced guilt when letting their loved ones go into respite care. Also, Winslow (2003) describes barriers to service use, such as care receiver resistance, reluctance of the carer, hassles for the carer, concerns over quality, and concerns over finances. Finally, the way in which the professional support is offered (intensity, frequency, flexibility, timing and location) does not always match the different individual needs (Koffman and Taylor, 1997; Gwyther, 1998; Coe and Neufeld, 1999; Perry and Bontinen, 2001; Clare, 2002). This often applies for standardized, and therefore non-flexible and non-personalized, care activities and programs. Biegel et al. (1993) found that people with dementia who do not use services or only in-home services were more functionally impaired and had inadequate informal support and more emotionally strained carers compared with users of out-of-home or both in- and out-of-home services. Coe and Neufeld (1999) describe the obstacles carers encounter when they have finally found the help they needed, “(. . .) for example, there was a delay in obtaining formal help, policies restricted the assistance that could be provided, and the application process was long and complex.” (p. 575).

The need for more flexible, personalized care and support is fully in line with the emphasis in the literature for many years that individual needs of patients and carers can considerably differ because of a number of personal and contextual factors, such as the symptoms of the person with dementia, carer characteristics and utilized coping strategies, the relationship between the carer and the person with dementia, the support systems available to the carer and the perception of the quality of the relief that is offered (Duijnste, 1992; Dröes et al., 1996; Dunkin and Anderson-Hanley, 1998; Burns, 2000; Proctor and Testad, 2005; Clare, 2002).

Some of the needs that people with dementia and their informal carers currently perceive as insufficiently met by regular care and support services might be alleviated, or even be met, using modern Information and Communication Technology (ICT). In this paper, we report on a review study that was aimed at getting insight into the state of the art of ICT solutions (in practice and research) that could contribute to meet these unmet needs of people with dementia and their informal carers. Based on the most frequently mentioned unmet needs by people with dementia and by their informal carers that were inventoried in field research among the target group (Dröes et al., 2006; Hancock et al., 2006; Roest et al., submitted), we focused our review study at the following need areas: (1) need for general and personalized information; (2) need for support with regard to symptoms of dementia; (3) need for social contact and company; and (4) need for health monitoring and perceived safety.

## 2. Method

### 2.1. Selection criteria

Within the four selected areas of needs a further subdivision into more specific themes was made (see Table 1) based on concrete needs mentioned by people with dementia in these areas, as described in the literature (Roest et al., 2007, submitted) and in three recent field studies that identified both met and unmet needs in people with dementia and their informal and formal carers. Dröes et al. (2006) inventoried general needs with respect to quality of life among people with dementia who attend meeting centres and regular day care centres ( $n = 106$ ) or who live in nursing homes ( $n = 37$ ). Hancock et al. (2006) researched unmet needs of people with dementia ( $n = 238$ ) in residential care using the Camberwell Assessment of Needs for the Elderly (CANE; Cummings et al., 1994). Roest et al. (submitted) conducted a large field study among 236 people with dementia living in the community and 322 carers using the Dutch version of the CANE (Dröes et al., 2004) and in-depth interviews.

From these studies it is clear that the need for general and personalized information mentioned by people with dementia refers to information on the diagnosis, condition, available support services (including information on support for example, memory problems) and care appointments. Informal carers want information on diagnosis, prognosis, treatment, care structures, day care facilities, and other services including legal and financial issues. The mentioned need for support with regard to symptoms of dementia refers to all types of instrumental support in daily life activities, including support to participate in activities and supervision/guidance.

The need for social contact and company refers specifically to ways of staying connected with family, friends and the social environment and to feel useful. Reported needs for health monitoring and perceived safety refer to the wish to be cared for and to feel secure when the disease progresses and disabilities increase.

For this review we focused on articles that were published before 15 November 2006 and met the following inclusion criteria:

- the article reports on people with cognitive disabilities related to dementia and/or their informal carers;
- the article discusses an ICT-device or application that has been tested within the target group;
- the ICT-device reported on in the article has proven to be helpful for the person with cognitive disabilities related to dementia and/or the informal carer on one of the selected needs areas of this review (see Table 1).

### 2.2. Search procedure

We searched the following databases: PubMed, PsychInfo, Cinahl, INSPEC and IEEE. Besides that we searched on the Internet using Google Scholar. Within these searches we used a list of key words for each needs area (see Table 2).

Table 1  
Studied needs areas and themes

Needs areas	Specific themes
1. Need for general and personalized information	A. Information on dementia B. Information on service offerings C. Information on legal and financial issues and on care and support services D. Information on personal condition, care appointments and care planning
2. Need for support with regard to symptoms of dementia	A. ICT compensation for disabilities such as memory problems in daily life activities B. ICT supporting the carer, flexibly and personalized, in providing instrumental care to the person with dementia C. ICT-support for people with dementia and carers regarding behavioral and psychological changes and how to cope with them D. Emotional support for patients and carers
3. The need for social contact and company for the person with dementia	Ways to stay connected with family, friends and the social environment and to be useful
4. The need for health monitoring and perceived safety for the person with dementia	The need to be cared for and to be safe when the disease progresses

Table 2  
Search strategy and keywords used in the search for ICT solutions in the different needs areas

Searched databases	General Keywords: Dementia[MeSH] OR dement*[ti] OR Alzheimer*[ti] Keywords in need areas and specific themes (A–D)	Information (A–D)	Support for symptoms (A–D)	Support in social contact	Health monitoring and safety
Pubmed PsychInfo Cinahl Google/ Google scholar		<p>AND Information[MeSH]<sup>a</sup></p> <p>(A) AND (diagnosis[MeSH] OR syndrome[MeSH] OR symptoms[tw]) AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[tw] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(B) AND (service[tw] OR Health service[MeSH] OR care[tw] OR support[tw] OR assistance[tw] OR help[tw] OR intervention[tw]) AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(C) AND (financial[tw] OR legal[MeSH] OR law[tw] OR costs[tw] OR benefits[tw]) AND (service[MeSH] OR Health service OR care[MeSH] OR support[MeSH] OR assistance[MeSH] OR help[tw] OR intervention[tw]) AND (electronic[MeSH] OR tele* OR technol*[MeSH] OR digital OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(D) AND (personal[MeSH] OR subjective[tw] OR individual[MeSH]) AND (condition[MeSH] OR status[MeSH] OR health[MeSH] OR severity[tw] OR stage[MeSH] OR “care plan”*[tw]) AND (e-health[tw] OR electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p>	<p>AND (service[MeSH] OR Health service[MeSH] OR care[MeSH] OR support[MeSH] OR assistance[MeSH] OR help[tw] OR intervention[tw])</p> <p>(A) AND (disabilities[tw] OR symptoms[tw] OR dysfunction*[MeSH] OR deficits[MeSH] OR problem[tw]) AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(B) AND (“informal carer”[tw] OR “informal caregiver” [tw] OR “family carer”[tw] OR “family caregiver[tw] OR caregiver” [MeSH] AND “instrumental care”[tw] AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR [MeSH] wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(C) AND coping[MeSH] AND/OR Adaptation[MeSH] AND (“person with dementia”[tw] OR patient[MeSH] OR “informal carer”[tw] OR caregiver[MeSH]) AND (“Behavioural symptoms”[tw] OR “Psychological symptoms”[tw] OR BPSD[tw] OR “psychosocial consequences”[tw]) AND (electronic[MeSH] OR Internet[MeSH] OR technological[MeSH] OR tele[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p> <p>(D) AND (emotion*[MeSH] OR affect*[MeSH] OR feeling*[MeSH]) AND (“person with dementia”[tw] OR patient[MeSH] OR “informal carer”[tw] OR “informal caregiver”[tw] OR “family carer”[tw] OR “family caregiver”[tw] OR caregiver[MeSH]) AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR software[MeSH] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer*[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw] OR robotic*[tw])</p>	<p>AND (service[MeSH] OR care[MeSH] OR support[tw] OR assistance[tw] OR help[tw] OR intervention[tw]) AND (“person with dementia”[tw] OR patient[MeSH] t) AND (social[MeSH] communication [MeSH] OR “social contact” OR company[tw] OR community[MeSH] OR group[tw] OR “social network” OR “social environment” OR “social support” OR interaction[MeSH] OR contact[tw] OR friend*[MeSH]) AND (electronic[MeSH] OR tele*[tw] OR technol*[MeSH] OR digital[tw] OR video*[tw] OR Internet[MeSH] OR website[tw] OR computer[MeSH] OR wireless[MeSH] OR phone[MeSH] OR personali*[tw] OR mobile[tw] OR pervasive[tw] OR ubiquitous[tw] OR domotic*[tw] OR “context-aware” [tw] OR sensors*[tw])</p>	<p>AND (health[MeSH] OR “life-signs”[wt] “health status”[tw] OR healthiness[MeSH] OR constitution[tw]) AND (“person with dementia” OR patient[MeSH]) AND (monitor*[MeSH] OR supervision[tw] OR attendance[tw] OR guidance[tw] OR control[tw] OR assistance[tw] OR care[MeSH] OR medicine[MeSH] OR pharmaco*[tw]) AND (safety[MeSH] OR security[tw] OR protection[MeSH] OR precaut*[tw] OR prevent*[MeSH]) AND (electronic[MeSH] OR detection[tw] OR device[MeSH] OR equipment and supplies[MeSH] OR prosthetic[tw] OR e-health[tw] OR technology[MeSH] OR tele*[MeSH] OR Bluetooth[tw] OR GPS[tw] OR biometric[MeSH] OR “alarm system” OR wireless[MeSH] OR locator[tw] OR application[tw] OR pervasive[tw] OR ubiquitous OR domotic* OR “context-aware” OR sensor[tw] OR gerontechnology[MeSH])</p>

<sup>a</sup> [MeSH] stands for several subcategories being subsumed under a keyword, such as dementia which comprises Alzheimer’s disease etc.; [tw] marks those words, that are best put in as text words, i.e. giving the best results; [ti] as used for dement\* or Alzheimer\* automatically searches for publications where these words occur in titles.

Table 3

Inventoried articles that discuss present ICT solutions in the area 'Need for general and personalized information' that have been proven helpful for people with cognitive disabilities related to dementia and for their carers (see also Table 1 for clarification of themes within need area)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
1. Need for general and personalized information (18 websites and 3 publications)				
Alzheimer Disease International	1A	People with dementia and informal carers	<a href="http://www.alz.co.uk">http://www.alz.co.uk</a> : Website with carer information and information for person with dementia	One thousand fifty visitors per month average in 2006
Alzheimer Europe	1A	Informal carers	<a href="http://www.alzheimer-europe.org">http://www.alzheimer-europe.org</a> : Website offering tips to informal carers; prevalence figures; research information	Circa 250,000 visitors of the website in 2005
Alzheimer Research Forum	1A	General public	<a href="http://www.alzforum.org">http://www.alzforum.org</a> : Website reports on the latest scientific findings, maintains public databases of essential research data and produces discussion forums <a href="http://www.aahsa.org">http://www.aahsa.org</a> : Digital social chart	Circa 2 million hits a year and 250,000 unique visits. About 75% of the registered users are researchers
American Association of Homes and services for the aging	1B	Elderly people		The members of the AAHSA serve two million people every day through mission-driven, not-for-profit organizations
American Department of Health and Human services	1B	General public	<a href="http://www.healthfinder.gov">http://www.healthfinder.gov</a> : Digital social chart, in addition online information on quality of care of services	Award-winning Federal website Since 1997 key resource for finding best government and non-profit health and human services information on the Internet. Received 63,647,896 hits in 2006
Belgian municipality	1B	General public	<a href="http://www.desocialekaart.be">http://www.desocialekaart.be</a> : Digital social chart	Contains concrete and up-to-date information on diverse welfare and healthcare services. No detailed information on number of hits or user characteristics available
Dutch Alzheimer Foundation	1A	People with dementia and (informal) carers.	e.g. <a href="http://www.alzheimer-nederland.nl">http://www.alzheimer-nederland.nl</a> : Digital information on: forms of dementia, practical, social and emotional consequences of dementia; online chat and forum <a href="http://www.denieuwezorgverzekering.nl">http://www.denieuwezorgverzekering.nl</a> : website for public information on health insurance and related developments; <a href="http://www.minvws.nl">http://www.minvws.nl</a> : information on the Exceptional Medical Expenses Act and the new Societal Support Act; <a href="http://www.zorgvoorbeter.nl">http://www.zorgvoorbeter.nl</a> : Website offering information on home care, quality of care, organizations, training and palliative care)	Eight hundred and fifty-four visitors per day 2006, 4000 signed up forum members 3 people with dementia actively participating in forum in January 2007
Dutch Ministry of Public Health, Welfare and Sports	1C	General public	<a href="http://www.kiesbeter.nl">http://www.kiesbeter.nl</a> : Digital social chart with service information for each province	<a href="http://www.denieuwezorgverzekering.nl">http://www.denieuwezorgverzekering.nl</a> , 2750 visitors per day average in 2006; <a href="http://www.minvws.nl">http://www.minvws.nl</a> , 168,400 visitors per month average in 2006; <a href="http://www.zorgvoorbeter.nl">http://www.zorgvoorbeter.nl</a> , 14,900 visitors per month average in 2006
Dutch Municipalities	1B	People with dementia and informal carers		Five thousands unique visitors per day average
Freeman et al. (2005)	1A	<i>N</i> = 5, early-stage Alzheimer's disease Single group, post-test design behavioral observational protocol and self-report satisfaction ratings	New website attuned to the needs and abilities of people with dementia. Comparison with existing website	Outcome on satisfaction ratings was used to set up recommendations regarding usability characteristics for websites to be used by people with dementia. Authors also composed a manual for designing websites for people with dementia

Table 3 (Continued)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
Stichting Vraagwijzer één loket	1B	General public	<a href="http://www.vraagwijzer.nl">http://www.vraagwijzer.nl</a> : Digital information on care, living and welfare	Vraagwijzer.nl is a national site that refers to local addresses where people can get help. Three thousand two hundred and thirty-five unique visitors in March 2007. 18 unique visitors per day average
Hassol et al. (2004)	1D	<i>N</i> = 4282, registered users of the application Online survey, focus groups with 25 users, individual interview with 10 primary care clinicians	Geisinger Health System, United States Mychart for patients, electronic communication with medical care providers, viewing selected parts of EHR, information on mental illnesses, services	Patients were satisfied that their medical information was complete and accurate. A minority of patients was mildly concerned about confidentiality and privacy of their information, and learning about abnormal test results electronically
The Henry Ford Hospital, Detroit	1D	50,000 patients of the Henry Ford Hospital, including people with dementia and carers.	MyHealth on <a href="http://www.henryford.com">http://www.henryford.com</a> : An application to view customized health information, obtain lab test results and renew prescriptions online	Largest patient portal in the USA with 50,000 patients using the application
Hulpjgids.nl, Private, non-profit initiative	1B	General public and professionals	<a href="http://www.hulpjgids.nl">http://www.hulpjgids.nl</a> : Online Information for professionals and public users on mental illnesses, therapies, medication, web chat. Extensive database of more than 1200 therapists (psychiatrists, psychotherapists, psychologists and alternative therapists), around 300 book titles and many links. The site also offers information on legal regulations and about what to do if one is dissatisfied with treatment or therapist. Problems can be discussed with a “web psychiatrist”	No information available on numbers of people of the website
Larner (2003)	1A	<i>N</i> = 104, with cognitive disorders of which 47 Alzheimer’s dementia, 11 Frontotemporal dementia, 3 Dementia with Lewy bodies, 3 Vascular dementia Survey, semi-structured interviews	Existing Internet websites and medical telephone helpline	Internet use by patients with cognitive disorders is common. The author endorses the view that the Internet may increase involvement of patients in supervising and documenting their own health care. Only 10% of all patients had used the medical telephone helpline
MEZZO, National Organization for informal carers and volunteer-carers	1B	Informal carers	<a href="http://www.mantelzorg.nl">http://www.mantelzorg.nl</a> : Website with regional information on care and support	Three hundred to five hundred unique visitors per day average
National Organization Regional Patient and Consumer platforms (LOREP)	1B	General public	<a href="http://www.zorgbelang-nederland.nl">http://www.zorgbelang-nederland.nl</a> : Digital social chart on health and welfare services Opportunity for feedback	No information available on numbers of people that visit the site
United States national Alzheimer Association	1A	(Informal) carers	<a href="http://www.alz.org">http://www.alz.org</a> : Website with information on dementia	Six million visitors annually. 4000 registered people with dementia or carers on the message boards in 2006
University of Florida	1A	(Informal) carers of people with dementia	<a href="http://www.alzonline.net">http://www.alzonline.net</a> : Specialized website with information for carers; Internet and telephone-based carer support; Alzheimer research Forum	Number of site visits in 2006: 184,628; number forum visits 2006: 6623; number of visits Spanish site 2006: 13,631

Table 4

Inventoried articles that discuss present ICT solutions in the area of ‘Need for support with regard to symptoms of dementia’ that haven been proven helpful for people with cognitive disabilities related to dementia and for their carers (see also Table 1 for clarification of themes within need area)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
2. Need for support with regard to symptoms of dementia (4 websites and 20 publications)				
Baruch et al. (2004)	2A	<i>N</i> = 1, patient with mild dementia living alone, case study	Computer system installed in bedroom and living room, giving automatic reminders, activity cues as well as showing time and date	Considerable reduction of support needed, reduction of night calls as well as reduction of perceived anxiety and confusion by person with dementia
Beauchamp et al. (2005)	2C	<i>N</i> = 299, carers in workforce. Randomized clinical trial	Multimedia support program, delivered via Internet	Improvement in carer depression, anxiety, level and frequency of stress, carer strain, carer self-efficacy, intention to seek help, perception of positive aspects of care giving
Van den Broek et al. (2000)	2A	<i>N</i> = 5, patients with prospective memory problems. Single group, pre-test–post-test design	Voice Organizer, electronic memory aid which can be ‘trained’ to recognize the user’s voice. The user can dictate a message and verbally set a date and time when the message should be repeated	Prospective memory improved with the Voice Organizer. Some patients showed improvement when they no longer used the device. This indicates a training effect
Camarinha-Matos and Asfaramesh (2004)	2B	Field assessment of prototype involving elderly and their relatives, care providers, decision-makers on social policy and software developers. Single group, post-test only	FP5-TeleCARE project. Aimed to design and develop a configurable framework focused on virtual communities for elderly support. The approach is based on the Internet and mobile agent technologies	The TeleCARE concept and its functionalities were well accepted by the potential users, although the system is at the research prototype stage and substantial engineering work is still necessary in order to make it a robust product
Dutch Alzheimer Foundation	2D	People with dementia and their carers	<a href="http://www.alzheimer-nederland.nl">http://www.alzheimer-nederland.nl</a> : Information on behavioral and psychological changes and how to cope, 24 h telephone service, contact fellow patients/carers	One thousand visitors per day average and 1,500 online orders for information materials in 2005
Gilliard and Hagen (2004)	2A	<i>N</i> = 80, people with mild to moderate dementia and informal carers, Single group, pre-test–post-test design	FP5-ENABLE project. Enabling technologies for people with dementia including Forget-Me-Not calendar Locator for lost objects Cooker usage monitor Medicine reminder	There is a potential for the devices assessed to support independence and reduce carers’ worry. A majority of the persons with dementia and their carers used the device, found it useful and were satisfied with the device. A majority of the persons reported good quality of life. Some of the devices reduced the general emotional burden of the carers
Glueckauf et al. (2004)	2C	<i>N</i> = 21, carers for persons with dementia, single group, pre-test–post-test design	ALZONLINE: Online tool to provide education and support services to carers of dementia web-and telephone- based care giving classes (hear, see and communicate with facilitator, slide presentations, communication with other carers) library: includes publications about dementia, frequently asked questions public message board links to federal, state and community resources	High utilization of message board, library and links. Enhancing use in broad population. Carer’s stress relieved and perceived self-efficacy feelings enhanced

Table 4 (Continued)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
Hofmann et al. (1996)	2A	<i>N</i> = 10, mild to moderate Alzheimer's single group, pre-test–post-test design	Interactive computer-based program, using photographs of the patient and his/her surroundings on a PC-touch screen to simulate an everyday task	Performing of computer-based training improved, showing that PwD are able to learn the use of technical programs and devices
Holte et al. (1998)	2A	<i>N</i> = 21, people with memory problems or cognitive impairment single group, post-test only	Forget-Me-Not device, electronic calendar showing date and day, can be personalized with photos indicating season or pictures of family	Proved usability helpful for orientation about day and date
Hulpgids.nl, Private, non-profit initiative	2C	General public	<a href="http://www.hulpgids.nl">http://www.hulpgids.nl</a> : Support for symptoms by consulting psychiatrists online, join forum, contact fellow patients/carers	No detailed information on number of hits or user characteristics available
Kinney et al. (2004)	2B	<i>N</i> = 38 (19 families) single group, post-test only	SAFE HOUSE PROJECT A web-based monitoring system, consisting of different technologies: A computer with Ethernet card, enabling Internet access Broadband service to access Internet Xanboo Smart Home Management System Cell phone with text messaging service	Carers burden in tracking/controlling loved one was relieved. Carers had more time to themselves. Relationships changed positively
Long and Williams (2005)	2C	Informal carers of people with dementia	<a href="http://www.james.psych.umn.edu/~grecc/caring.htm">http://www.james.psych.umn.edu/~grecc/caring.htm</a> : Online pamphlets for carers with information on bureaucratic issues, cope with decline, hearing and vision problems and bowel and urinary incontinence	No detailed information on number of hits or user characteristics available
Magnusson et al. (2002)	2B	Long-term users of the system, 26% dementia, informal carers Single group, post-test only	FP4- ACTION project. Assisting Carers using Telematics Interventions to meet Older Persons' Needs	Effective informal social support networks via videophone for family carers. High levels of global usability. The system enabled enriching activities to be carried out by carers and the persons they care for. Early access to the system would help to develop the novice carer's caring competence
Oriani et al. (2003)	2A	<i>N</i> = 5, outpatients with Alzheimer's Disease, three experimental conditions single group, post-test only	Electronic memory aid (EMA) consisting of a vocally recorded appointment or task, to be pre-programmed by date and time. At given date audio-alarm is activated to which the patient has to react by pressing a button	Use of EMA improved prospective memory. Support via EMA was more useful than written list or free recall
Poon et al. (2005)	2A	<i>N</i> = 22, mild dementia or mild cognitive impairment. Clinical trial	Comparison of cognitive intervention delivered face to face or via a video-conferencing system. Cognitive intervention comprised assessment, intervention and psychosocial support	With regard to attention, memory and language areas, Telemedicine in comparison to face-to-face treatment as effective, acceptable and feasible Positive evaluation by patients because of it is cost- and time saving character Telemedicine improves accessibility of support services in communities
Szymkowiak et al. (2004)	2A	<i>N</i> = 8 (two trials, four patients each) single group, post-test only	PDA memory aid system: automatic reminders, action prompts and calendar function	Positive evaluation of usability and functionality of system; Recommendations for improvement of PDA's; decrease of worry in carers

Tarraga et al. (2006)	2A	<i>N</i> = 46, people with Alzheimer's disease. Randomized clinical trial	Interactive multimedia tool of cognitive stimulation, IMIS (15 difficulty levels; 19 exercises across different domains)	IMIS improved MMSE and ADAS-Cog scores, improvement was maintained over 24 weeks of duration of program; IMIS improvement significantly higher than in the group that received only integrated psycho-social stimulation IPPS (daily program in day care centres)
Tunstall	2B	People with dementia	Tunstall telecare system <a href="http://www.tunstalltown.com">http://www.tunstalltown.com</a> Telecare devices such as doorbells, videophones, home response systems, detectors tailored to individual needs	Devices are used in studies
Vehvilainen et al. (2002)	2D	<i>N</i> = 42, patient-carer dyads, Alzheimer's Disease. Field trial, post-test only	A decision-making and emotional support platform for carers of people with Alzheimer's Disease	Positive results showed that over three quarters of the test group would continue to use the service on a daily basis
White and Dorman (2000)	2C	<i>N</i> = 1015, mailing list; 178 different users actively posted messages. Content analysis of website postings	Alzheimer mailing list, created by Washington University Alzheimer's Disease Research Center (ADRC) <a href="http://www.adrc.wustl.edu/Alzheimer">http://www.adrc.wustl.edu/Alzheimer</a>	Content analysis revealed that mailing list provided participants with information, emotional support and advice
Wilson et al. (2001)	2A	<i>N</i> = 143, 7.6% of participants diagnosed with an acquired progressive brain injury (e.g. Alzheimer's Disease). Randomized clinical trial	NeuroPage, prosthetic memory and activity cueing. The system involves an arrangement of microcomputers linked to a conventional computer memory and, by modem, to a paging system.	Significant reduction of everyday memory and planning failures Improvement on carrying out daily activities
Woolham (2005)	2B	<i>N</i> = 233, people with dementia; <i>N</i> = 173, control group. Pre-test–post-test matched control group design	Safe at home project: assistive and telecare technology device installed at home	Compared to control group, group with technology had fewer care visits, lived longer in community, improved confidence, decreased carer concerns
Yasuda et al. (2006)	2C	<i>N</i> = 3, mild to moderate dementia. Alzheimer's Disease, vascular dementia, mixed AD-VD. Case studies, pre-test-post-test design	Music and messages automatically output by IC recorder for guiding behavior in people with dementia	Automatically output music and messages were highly effective for guiding person with dementia to go to a day care centre, behave more peacefully and eat more meals. Method could easily be applied to other activities of daily life
Zanetti et al. (2000)	2A	<i>N</i> = 5, people with Alzheimer's disease. Pre-test–post-test control group design	Electronic agenda, a memory aid to support people with dementia in prospective memory tasks	Patients showed significant improvements on several prospective memory tasks. Two out of five patients obtained a maximum score on the tasks when using the device

Table 5

Inventoried articles that discuss present ICT solutions in the area of ‘Need for social contact and company for the person with dementia’ that haven been proven helpful for people with cognitive disabilities related to dementia and for their carers (see also Table 1 for clarification of themes within need area)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
3. Need for social contact and company for the person with dementia (11 publications)				
Ager and Aalykke (2001)	3	<i>N</i> = 18, representatives of user groups. Intellectual disability, acquired brain injury and dementia. Single group, post-test only	FP4-TASC project, Telematics Applications Supporting Cognition. A flexible, modular microcomputer support system for persons with cognitive disabilities. The five modules comprising the TASC software are The Planner, for planning and time management The Communicator, for communication by mobile The Prompter, for guidance The Supervisor, for environmental control The Information provider, for information provision	Increased independence. Self-assurance and improved social contacts
Alm et al. (2003)	3	<i>N</i> = 18, patient/carer dyads. Single group, post-test only	A multimedia-based reminiscence tool to help improve communication and social interaction of people with dementia	Increased social interaction as a result of enhanced and prolonged conversations
Bourgeois et al. (2001)	3	<i>N</i> = 66, people with dementia; treatment and control group (33). Randomized clinical trial	Memory aid to improve communication/ social skills	Increased quality of life, enhanced information sharing and social closeness
Gilliard and Hagen (2004)	3	<i>N</i> = 80, people with mild to moderate dementia and informal carers. Single group, pre-test–post-test design	FP5-ENABLE project. Enabling technologies for people with dementia including Picture gramophone My history device Pre-programmable telephone	Assistive technologies can facilitate independent living of people with dementia and promote their well-being by giving positive experiences. A majority of the persons with dementia and their carers used the device, found it useful and were satisfied with the device. A majority of the persons reported good quality of life. Some devices reduced the general emotional burden of the carers
Kort (2005)	3	<i>N</i> = 4, people with Alzheimer’s dementia. Single group, pre-test–post-test design	Mobile Telecoach Mobile phone with just one button, answering incoming calls automatically	Positive effects on self-esteem. Positive social experiences
Lekeu et al. (2002)	3	<i>N</i> = 2, people with dementia Single group, pre-test–post-test design	Training program consisting of spaced-retrieval and errorless learning techniques, to teach use of mobile phone	Effective in teaching how to use a mobile phone
Mickus and Luz (2002)	3	<i>N</i> = 20, patient carer dyads in nursing home. Single group, post-test only	Videophones, used for televisits	Usability of videophones, Increased number of social interactions
Olsen et al. (2000)	3	<i>N</i> = 20, 75% people with dementia participating in a social day care program. Single group, pre-test–post-test design	Music Memory Lane (MML) and Video Memory Lane (VML) present nostalgic music and videos to people with Alzheimer’s Disease	MML and VML have a favorable impact on engagement, stimulated positive affect and activity-related talking, while also reducing fidgeting

Sakairi (2004)	3	<i>N</i> = 8, patients with senile dementia. Single group, pre-test–post-test design	A robot dog (AIBO) used to stimulate communication	With the introduction of the toy dog patient activity and communication between staff and other users was improved
Savenstedt et al. (2003)	3	<i>N</i> = 7, patient-family dyads. Single group, post-test only	A video phone system used to communicate with family members	More frequent contacts as compared to usual physical meetings
Tamura et al. (2004)	3	<i>N</i> = 13, severe dementia. Single group, post-test only	Entertainment robot (AIBO) versus motorized toy dog	Both AIBO and the toy dog effectively increased patient activity during occupational therapy. Results indicate increased socialization and social activity in the presence of AIBO and toy dog

---

Table 6

Inventoried articles that discuss present ICT solutions in the area of ‘Need for health monitoring and perceived safety for the person with dementia’ that haven been proven helpful for people with cognitive disabilities related to dementia and for their carers (See also Table 1 for clarification of themes within need area)

Need area/author(s)	Theme	Research sample/target group and design	Device/application	Outcome/results
4. Need for health monitoring and perceived safety (12 publications)				
Algase et al. (2003)	4	<i>N</i> = 178, dementia diagnosed by DSM-IV criteria Single group, post-test only	Biomechanical activity devices for capturing movement which provide a way to index wandering. (Actillum, StepWatch, Step Sensor and TriTrac-R3D)	The StepWatch was superior. This device may be useful in assessing the amount and daily distribution of wandering behavior
Blackburn (1988)	4	<i>N</i> = 77, elderly mentally infirm patients in district general hospital. Baseline measurement ( <i>N</i> = 29). Measurement after installation of the system ( <i>N</i> = 48). Number of nursing staff not reported on. Cross-sectional design for patients. Single group, pre-test–post-test design for nursing staff	Monitoring device attached to patients’ clothes with boundary alarms. Buzzers and lights at strategic points in the ward	Reduction in potentially serious incidents and staff anxiety levels and a more effective use of nursing time
Chan et al. (2002)	4	<i>N</i> = 1, moderate dementia. Case study, post-test only	Smart tool for the assessment of the mobility of patients with motor disorders. System consists of a network of IR sensors installed in patient’s housing in a long-stay setting	The system is valid in providing the carer with data about real nocturnal activity and shows great potential for providing data concerning current mobility that may be indicative of ongoing changes in people with dementia
Lin et al. (2006)	4	<i>N</i> = 11, healthy individuals. Single group, post-test only	A comprehensive stray prevention system to facilitate minimal intrusion on patient lifestyle	Reviewed by 11 individuals with positive feedback on suitability for application in searching for elderly patients with dementia
Ikarashi et al. (2002)	4	Patient with dementia and family members. Case study, post-test only	A GPS location monitoring system	Reduced mental stress of family members
Masuda (2002)	4	<i>N</i> = 3, dementia patients. Single group, post-test only	A bedside monitor to detect wandering at night	Successfully detected 30 cases of wandering during which zero falls or injuries were recorded
Miskelly (2004)	4	<i>N</i> = 12, people with dementia, 5 in hospital setting, 4 in residential homes, 3 living at home. Single group, post-test only	Electronic tracking system (bracelet and monitoring station)	Reliably and successfully detects external wandering. Good compliance in patients
Miskelly (2005)	4	<i>N</i> = 11, Patients with dementia in nursing home, carried device in either shoulder bag or waist bag. Single group, post-test only	Mobile phone with GPS used for electronic tracking	Successful localization of lost people within a range of 5 m
Ogawa et al. (2004)	4	<i>N</i> = 2, a wandering elderly person and carer. Single group, post-test only	Mobile PHS pendant using radio waves for location tracking	Successful location of wandering person at 1 min intervals at a range of 60 m, tested at 15 different locations
Orpwood et al. (2005)	4	<i>N</i> = 138, carers of people with dementia. 45 from Alzheimer’s Society day clubs, 45 from local authority centres and 48 involved in focus groups by mental health team. <i>N</i> = 30, carers in survey to prioritize preferred ICT solution	Smart home environment with different services, e.g. a lighting system during the night, reminding system, cooking monitor	Working closely with personal carers of people with dementia ensures that the interaction between individuals with dementia and assistive technology will be more effective
Paavilainen et al. (2005)	4	<i>N</i> = 42, non-demented and mild to severe dementia patients in nursing homes Between group design, post-test only	IST Vivago <sup>®</sup> WristCare, an active social alarm which provides continuous telemetric monitoring of the user’s circadian activity	Telemetric actigraphy: useful method in long-term screening and follow-up in elderly subjects for sleep- and circadian rhythm-related problems associated with dementia
Shimizu et al. (2000)	4	<i>N</i> = 2, a wandering elderly person and carer. Single group, post-test only	GPS location based service for people with dementia	Successful location tracking in adverse weather conditions and in close proximity to tall buildings

The snowball effect furthermore enabled us to retrieve other publications based on the references in the publications found initially. The search procedure did not reveal information derived from unpublished studies.

### 2.3. Analysis

The searches within the mentioned databases initially resulted in 165 publications and websites. The analysis was performed by two groups of researchers; the first group consisted of three people with expertise in the clinical domain (AR, SL, RMD); the second group consisted of two people with expertise in information and communication technology (RJD, MDM). All articles were independently assessed with regard to the inclusion criteria by two researchers and compared afterwards. In case of discrepancies the articles were considered again until consensus was reached. This resulted in 22 (examples of) relevant websites and 46 articles that satisfied the inclusion criteria for this review (see Tables 3–6). The selected articles were analyzed with regard to needs area, research sample and research design, functionality of the ICT solution/device and outcome or results for the target group (see Tables 3–6). The content of the selected ICT solutions and devices is described briefly in the Section 3.

## 3. Results

### 3.1. Need for general and personalized information

In this needs area 18 relevant websites and three publications were found that satisfied the inclusion criteria.

#### 3.1.1. Information on dementia

Larner (2003) demonstrated that the use of the Internet by people with cognitive disorders and their carers is common and may increase involvement of patients in supervising and documenting their own healthcare.

The websites of Alzheimer associations provide digital information on dementia (e.g. <http://www.alzheimer-europe.org> or <http://www.alz.org>). Apart from prevalence figures, research information and advice on the practical, social and emotional consequences of the disease, some of these sites offer the opportunity to chat with fellow patients and carers (e.g. <http://www.alzheimer-nederland.nl>). A few sites include an informative section for people with dementia themselves (e.g. <http://www.alz.co.uk>).

As part of the UK's Alzheimer's Society project 'Learning to Live with Dementia', Freeman et al. (2005) set up guidelines for accessible writing and web design for people with mild to moderate dementia. In order to attune (online) text to the needs of people with dementia they designed and evaluated a website in close collaboration with the target-users, which improved the accessibility of the website.

The University of Florida provides a specialized website (<http://www.alzonline.net>) where carers are offered information on dementia as well as Internet- and telephone-based support and education. The Alzheimer Research Forum (<http://www.alzforum.org>) includes detailed information on clinical guidelines, the medical workup, drugs and non-pharmacological therapies for health care professionals, researchers and lay people.

#### 3.1.2. Information on service offerings

Besides printed guides containing an overview of healthcare and welfare services, recently online 'social charts' have become available in several European countries. They provide detailed information on national or regional services for professionals and/or specific patient groups. Social charts are offered by different organizations and therefore vary with regard to target-users, ease-of-use and completeness. Some popular Dutch social charts are provided by the National Institute for public health and environment (RIVM: <http://www.kiesbeter.nl>), the Trade organization of regional consumer- and patient interest organizations (<http://www.zorgbelang-nederland.nl>) and the National society of informal- and voluntary carers (MEZZO; <http://www.mantelzorg.nl>). The site <http://www.hulpguides.nl> focuses on mental health care and contains a database of over 1200 clinicians, therapists and psychiatrists. More general sites (e.g. <http://www.vraagwijzer.nl> and <http://www.2zw-adreswijzer.nl>) offer information on healthcare, welfare, housing, social security, income, education and policy to a broad audience.

Social charts in other countries, such as <http://www.desocialekaart.be> in Belgium, <http://www.alzonline.net> in the state of Florida and <http://www.aahsa.org> in the USA are in essence not very different, though some, like <http://www.healthfinder.gov>, also provide visitors with comprehensive information on the quality of care.

The social charts discussed above are typically quite static, generic (not specifically intended for people with dementia or their carers) and mainly provide an, often incomplete, list of addresses for services in the region. Personalized, context-sensitive and demand-orientated sites which exist for other diagnostics groups (e.g. <http://www.voorlichtingopmaat.nl> for rheumatism patients or <http://www.stoppain.org> for pain patients) are lacking for people with dementia and their informal carers.

### 3.1.3. Information on legal and financial issues and on care and support services

The websites of ministries responsible for public health, like the Dutch ministry of Public health, Welfare and Sports (VWS) offer information on financial issues, legislation, health insurance and other subjects related to financial and legal developments in care and support services (<http://www.minvws.nl> or <http://www.denieuwezorgverzekering.nl>). With the website <http://www.zorgvoorbeter.nl>, which is part of a national action program, the Dutch ministry of VWS aims to improve quality of care in home care and institutional care for the elderly and disabled by providing transparency, quality indicators and stimulating improvement trajectories in healthcare organizations. The site also offers descriptions and examples of high-quality care for people with dementia. A telephone service (the ‘Mantelzorglijn’) offers support to informal carers and can also be contacted for legal and financial advice with regard to care and support services.

### 3.1.4. Information on personal condition, care appointments and care planning

In the USA, the Geisinger Health System with its ‘MyChart’ application allows patients to communicate electronically with their medical care providers and view selected portions of their Electronic Health Record (EHR) (Hassol et al., 2004). Likewise, the Henry Ford Hospital in Detroit, Michigan has over 50,000 patients using an application called ‘MyHealth’ on [henryford.com](http://www.henryford.com) to view customized health information, obtain lab test results and renew prescriptions online (<http://www.henryford.com>).

Recently, a new system called the Patient Access Electronic Record System (PAERS) was trailed in the UK, which allows patients visiting their General Practitioner (GP) to look up their health records and book in for their appointments (Honeyman et al., 2005). It is not reported whether people with dementia or their informal carers use this system.

Although EHR’s are applied on a small scale in hospitals, home care organizations and GP networks in many European countries, patients and informal carers have no access to these records yet. As an increasing number of governments demand that an Electronic Medical Dossier (EMD) and, at a later stage, an Electronic Patient Dossier (EPD) needs to be available to every party involved, organizations like the Dutch NICTIZ (<http://www.nictiz.nl>) work on standardizing information exchange between healthcare providers. An important emerging standard to exchange basic medical information is the HL7v3 standard. Apart from memory aids (see next paragraph) ICT solutions that inform people with dementia on their personal condition, appointments and care planning are still in the research and development stage.

## 3.2. Need for support with regard to symptoms of dementia

In the needs area support with regard to symptoms of dementia we included four websites and 20 publications.

### 3.2.1. ICT compensation for disabilities such as memory problems

To help people with dementia undertake specific types of time-linked tasks (e.g. taking medication, keeping appointments or following a daily schedule) various possibilities of automatic reminders via Electronic Memory Aids (EMAs) are explored (e.g. Kim et al., 2000; Wilson et al., 2001; Hart et al., 2004). Kim et al. (2000) studied the use of a palmtop computer based memory aid by persons with functional impairments of prospective memory and found this technology to be useful in a high proportion of patients for assisting memory dependent functions. Hart et al. (2004) conducted a consumer survey study to ascertain experiences and attitudes regarding the use of portable electronic devices as memory and organizational aids. They concluded that these devices are acceptable or desirable for use as compensatory aids by consumers with moderate to severe brain injury. While these initiatives differ in approach, they agree on the usability of memory aids for persons with dementia, the importance of defining their own reminders and of giving them and their informal carers control over the reminders (e.g. Wilson et al., 2001; Inglis et al., 2003). Most researchers recognize the importance of involving informal carers in the implementation of EMAs as carers motivate

the person with dementia to use the EMA but also benefit themselves, as use of an EMA by a person with dementia reduces feelings of worry in the informal carer (Szymkowiak et al., 2004).

Only a few studies have investigated the effect of EMAs for people with dementia (Van den Broek et al., 2000; Zanetti et al., 2000; Wilson et al., 2001.). Most studies are uncontrolled and based on a small sample, but justify hope for future development. Van den Broek et al. (2000) studied the use of the Voice Organizer, a device on which a message can be recorded by the user and that plays it back on a specified date and time. In the patients with memory problems that participated in the study the prospective memory improved directly after the device was taken into use.

Zanetti et al. (2000) asked five patients with Alzheimer's Disease to conduct seven prospective memory tasks on set times with the help of an electronic agenda and compared the effort with a control condition in which the same tasks had to be performed without the electronic agenda. All participants performed significantly better on the memory tasks when using the electronic agenda. Wilson et al. (2001) report on the effectiveness and usability of the 'NeuroPage' system (Hersh and Treadgold, 1994), a pager system which allows for a large number of reminders and prompts for various activities (<http://www.neuropage.nhs.uk>). Within a mixed diagnostic group with memory disorders, including people with dementia, NeuroPage as well as a pocket-computer memory enhancement system were evaluated and found to be workable although motivation and remembering to enter relevant information did limit overall usefulness (Wilson et al., 2001). EMAs appear particularly suited to support prospective memory and, although they are generally more expensive than non-electronic memory aids, require little training (see review Fritschy et al., 2004).

An electronic calendar to aid persons with memory problems or cognitive impairment, called the Forget-Me-Not device, was produced within the European Technology, Ethics and Dementia (TED) project and could be useful for persons who are confused about day and date (Holte et al., 1998). An EMA informing the person verbally, using vocal recordings, of specific appointments or tasks, also showed positive effects (Oriani et al., 2003). Inglis et al. (2003) and Szymkowiak et al. (2004) utilized a Personal Digital Assistant (PDA) to remind users of daily tasks, telephone numbers, appointments and birthdays. Users appreciated the device, though some visually or hearing impaired users preferred a reminder in the form of vibration.

Baruch et al. (2004) observed beneficial effects of computer systems on orientation, feelings of anxiety and independency in a patient suffering from Alzheimer's disease. Computer screens in the bedroom and living room reduced the needed support and the number of nighttime calls to the informal carer.

Within the multi-national ENABLE project (Enabling technologies for people with dementia) twelve existing products (-concepts) were identified and their effectiveness to support memory, improve quality of life and reduce the carer burden was assessed. These assistive technologies were found to facilitate independent living and some devices may reduce anxiety in people with dementia as well as their informal carers (Gilliard and Hagen, 2004). EMAs, sometimes combined with interventions like visual imagery, errorless learning or spaced retrieval techniques, proved efficacious in stimulating memory in people with dementia (see review Grandmaison and Simard, 2003).

Cognitive intervention is another approach to support people with dementia in their memory function and can be presented utilizing video conference, computer-based, or Internet-based systems.

In comparing cognitive intervention via a video conference system with face-to-face intervention, Poon et al. (2005) found telemedicine to be feasible, effective on memory, language and attention and highly accepted in people with dementia. Hofmann et al. (1996) used an interactive computer-based program to train people with dementia to use a touch screen and showed improved performance in computer-program tasks. To determine the usefulness of an Interactive Multimedia Internet-based System (IMIS) in people with suspected Alzheimer's disease, Tarraga et al. (2006) compared IMIS in combination with an Integrated Psycho-stimulation Program (IPP) and pharmacological treatment to IPP combined with pharmacological treatment and pharmacological treatment alone. They found both IMIS and IPP to improve cognition and the IMIS program to provide improvement above that seen with IPP alone.

### 3.2.2. ICT supporting the carer, flexibly and personalized, in instrumental care

Formal and informal carers in a number of European and North-American countries utilize forms of telemedicine such as video conferencing, telemonitoring and telecare to create more capacity by reducing travel time and increase quality of care by allowing more frequent contact and quick referrals to a specialist. Several projects aimed to develop telecare home services that enable persons with dementia to live independently and offer support to their informal carers.

The European 4th Framework Program (FP4) project ACTION (Assisting Carers using Telematics Interventions to meet Older persons Needs) focused on the empowerment of family carers to help maintain autonomy, independence

and quality of life in frail elderly. Familiar electronic equipment like TV and telephone was combined with modern ICT to improve carers' coping skills in daily care and emergency interventions and to offer financial information and emotional support (Magnusson et al., 1998a,b). A Swedish contribution to the project comprised a videophone to facilitate communication between healthcare providers and patients and their families. Magnusson et al. (2002) found that families were quicker to request information, education and support from professional carers and valued the informal support network of family carers to share experiences.

The FP5 project TeleCARE aimed to design and develop a flexible infrastructure of remote supervision and assistance services to facilitate independent lifestyles and to improve quality of life, confidence, well-being and safety in elderly people, including people with dementia. Partial validation was achieved with field assessments involving four classes of potential users including elderly and their relatives and healthcare providers. However, field-tests for fine tuning and acceptance of the technology still need to be carried out (Camarinha-Matos and Asfaramesh, 2004). Another telecare project is the 'Tunstall telecare system', which can be tailored to suit the needs of people with dementia and protect them from dangers such as wandering, fires and floods (<http://www.tunstalltown.com>).

Within the 'Safe-at-home' project, Woolham (2005) conducted a large-scale study into the effect of different items of assistive and telecare technology. An intervention-group of 233 people with dementia was compared to a matched control group of 173. After intervention the two groups differed significantly in the number of services received, visits and contact-hours per week, resulting in lower costs in the intervention-group. The technology was found reliable and a majority of carers reported reduced feelings of concern for a person's safety. Almost half of the carers felt the project had improved the confidence of the user in their ability to look after themselves safely.

The experiences of 19 informal carers with a web-based home monitoring system, consisting of broadband Internet access, an Ethernet card, the Xanboo Smart Home Management system and a cell phone with text messaging, were explored by Kinney et al. (2004) in the SAFE house project. Carers received training on how to use the system prior to installation and were able to use the system during the intervention period of 24 weeks. Researchers reported reduced carer burden. Cell phone alerts facilitated keeping track of loved ones and as relatives called more often, relationships improved (Kinney et al., 2004).

### 3.2.3. *ICT-support for people with dementia and carers with regard to behavioral and psychological changes and how to cope with them*

Information on behavioral and psychological changes in people with dementia, how to cope with them and where to find support can be found on websites of Alzheimer societies (e.g. <http://www.alzheimer-nederland.nl>). Alzheimer Nederland also offers a 24-h telephone service for information, support and advice. The Dutch site <http://www.hulp.gids.nl> offers psychiatric consultation, forum discussions and fellow-patient contact. On <http://www.vraagwijzer.nl> one can get help clarifying a healthcare demand, get advice on adequate care and support in how to receive the desired type of care.

The Geriatric Research, Education and Clinical Centre (GRECC) of the Minneapolis Department of Veteran Affairs Medical Centre offers practical online information on a number of topics related to caring for a patient with dementia (e.g. how to handle a loved one's decline, how to cope with bowel and urinary incontinence) (<http://www.james.psych.umn.edu/~grecc/caring.htm>; Long and Williams, 2005). The US website <http://www.alzonline.net> provides Internet- and telephone-based support and education for people with Alzheimer's disease and other dementias as well as support for their carers. Utilizing training videos and downloadable information, the site offers classes to address a wide range of topics related to dementia care (e.g. medication management, dealing with the stress of caring and performing daily tasks). An initial program evaluation showed reasonable effectiveness of AlzOnline's Positive Caregiving Classes. Carer self-efficacy improved and subjective carer burden was reduced but positive dimensions of the care giving experience and perceptions of time-burden in providing care were unaltered (Glueckauf et al., 2004). The efficacy of 'Care-giver's Friend', an Internet-based multimedia support program designed to support family carers employed in the workforce, was tested by Beauchamp et al. (2005) in 299 participants using a pre-test/post-test design. Results showed significant benefits on carer depression and anxiety and improvements in perceived stress, strain and positive aspects of care giving.

Guiding the behavior (e.g. to go to a daycare centre, eat more meals) of people with dementia by music and sung messages output by an IC recorder was found to be highly effective (Yasuda et al., 2006). The researchers suggest that music can have a strong effect on mental stability and the method can easily be applied to other activities of daily life.

### 3.2.4. Emotional support for people with dementia and carers

White and Dorman (2000) analyzed the contents of messages posted on an Alzheimer mail group and concluded that the opportunity to share, unburden or vent, is perceived as empowering and helpful to carers. Almost every Alzheimer association offers a website where people with dementia and their carers can join forums, post-messages or chat with fellow sufferers (e.g. <http://www.alzheimer-nederland.nl>, <http://www.pick.nl>). Many Alzheimer associations also have a 24-h telephone support service for emotional support and information on regional support services like Alzheimer café's and meeting centres.

An Internet-based application called Alzheimer's Carer Internet Support System (ACISS), designed to provide carers with clinical, decision-making and emotional support, was evaluated in a 6-month field trial of 42 carer/patient dyads. Preliminary results showed the system to be beneficial to carers of people with dementia (Vehvilainen et al., 2002).

### 3.3. Need for social contact and company for the person with dementia

In the area need for social contact and company 11 publications were included. Simplified mobile phones for elderly persons to contact informal carers or relatives directly have recently been developed. Lekeu et al. (2002) demonstrated that people with mild dementia are able to learn how to use a mobile phone with an 'errorless learning'-method (Clare et al., 2000). The 'Mobile Telecoach', a one-button mobile phone which allows direct answering, had an effect on positive social experiences and self-esteem in people with dementia. However, users were not satisfied with its size, weight and battery life (Kort, 2005). The FP4 project MORE was aimed at redesigning existing mobile phones and simplifying the user interface to meet the many differing needs of elderly and disabled people. Various MORE-based telephones with an integral GPS function were produced by Benefon, but none of these have come into use as navigation support for people with disabilities (Lindström, 2005).

Within the FP4 project Telematic Applications Supporting Cognition (TASC), five software-modules were developed for supporting persons with developmental disability, brain injury or dementia in information provision, communication, environmental control, planning and time management. The TASC-system runs on a standard computer; the modules can be used separately or together, are designed to be compatible with future extensions and can be programmed to fit the needs of an individual user. Usability and functional effectiveness of alpha-versions of the system were trailed with representatives of user-groups. The amended system was implemented with 18 users across five validation sites. The field trials indicate that the TASC-system can reduce the need for help from others with daily activities, leading to increased independence and quality of life and reduced costs for assistance and institutional care (Ager and Aalykke, 2001). The ENABLE project included a number of devices to provide pleasure or comfort and facilitate communication such as a picture gramophone (a do-it-yourself multimedia program), a 'My history' device (PC with touch screen showing pictures of people and places with familiar voice narrative) and a pre-programmable telephone (calls initiated by just pressing one large button containing a name or photo). Trials among people with mild to moderate dementia indicate that these assistive technologies can enhance well-being by giving positive experiences, and reduce anxiety in people with dementia and their informal carers (Gilliard and Hagen, 2004).

The 'Musical Memory Lane' built in a 1930s Philco radio cabinet and the 'Video Memory Lane' housed in a 1950s television cabinet, present nostalgic music and videos to people with Alzheimer's Disease in an easy-to-access, push-button, picture format. Systematic observations indicate that the Memory Lanes have a favorable impact on engagement, stimulate positive affect and activity-related talking, while also reducing fidgeting (Olsen et al., 2000).

'Televisits', contact between elderly people living in a nursing home and their family via a videophone, was demonstrated to promote social contact by Mickus and Luz (2002). In a similar study, Savenstedt et al. (2003) showed that a videophone as a means of communication between patient and family reduced feelings of guilt in family members, allowed more frequent visits than was possible with face-to-face visits, and let family members see the physical and emotional state of the patient on a daily basis. In some cases the conversations were more focused and of better quality than during face-to-face visits. In most cases, however, more emphasis was placed on the family member to direct and lead the conversations which was seen by many as demanding. The relationship between staff and family members improved as a side effect of staff helping the patient use the videophone.

Tamura et al. (2004) found that a motorized toy dog can effectively reduce wandering and agitation after dinner in people with dementia. The robot-dog AIBO as well as a motorized toy dog proved to be effective in increasing patient activity and spontaneous speech during occupational therapy. Sakairi (2004) showed that the introduction of AIBO

increased the number of utterances in people with dementia. These results indicate that socialization and social activity can increase in the presence of a toy dog and AIBO. Alm et al. (2003) suggest that with the use of multimedia as a source of reminiscence, patients exhibit more control of the direction of the conversation. They developed a touch-screen display to convey photographs, video or music, and compared it to traditional reminiscence methods. The patients using the multimedia system showed stronger and more prolonged engagement during the conversations.

#### 3.4. Need for health monitoring and perceived safety

Implementing monitoring technologies and detection devices or alarm systems inside and outside the home of elderly persons is potentially useful to enhance (perceived) safety and security of the person suffering from dementia as well as carers. Over the last few decades, detection devices and alarm systems for health problems and safety have been developed for different diagnostic groups, but we found only 12 studies that investigated the effect of these devices among people with dementia.

In a study comparing four biomechanical activity devices to index wandering, Algate et al. (2003) found the Step Watch particularly able to assess amount and daily course of wandering behavior in people with dementia. In this context, Boundary alarms (activated by a wristband) or electronic tagging with bracelets and monitoring stations were found to be effective, reliable and successful in detecting wandering as well as reducing patient and carer stress (Blackburn, 1988; Miskelly, 2004). A bedside monitoring system tested in a hospital setting with three patients with dementia who frequently wandered during the night, additionally provides floor lighting upon wandering detection and relays an alarm to a personal handheld device alerting the carer to the situation so that the necessary intervention can be performed (Masuda et al., 2002). A straying prevention system developed by Lin et al. (2006) can counteract wandering without interfering with the daily life activities of the person suffering from dementia. The system consists of indoor residence monitoring, outdoor activity monitoring, emergency rescue service and remote monitoring that can be accessed via a number of mobile devices such as a mobile phone, PDA, notebook or computer. The indoor monitor detects movement between certain areas within the home. Outdoor activities, accompanied or unaccompanied, are monitored in a pre-set activity area, which is activated by pressing a button on the location tracking device. Messages, or alarms respectively, are generated and forwarded via mobile communication (GSM) if the patient leaves the home environment or the activity area. In case of an emergency, the person with dementia can activate an emergency button which sends a message to the call centre where the situation, location and geographical information and location coordinates are analyzed and relayed to care providers, search teams and family members. Conversely, through the secure remote monitoring facility included in the system, family members or care providers can access patient's location information at any time by logging onto the system. Although it was not tested with patients of the target group, the system was expected to be beneficial in helping locate elderly people with dementia by 11 volunteers who reviewed the system. Paavilainen et al. (2005) tested an active social alarm system, IST Vivago<sup>®</sup> WristCare, which provides continuous telemetric monitoring of the user's activity. Results of this study support the use of telemetric actigraphy in long-term screening and follow-up of elderly subjects for sleep and circadian rhythm-related problems associated with dementia, and changes in functional capacity.

A fully automatic multi-sensor system, composed of infra red (IR) sensors connected to a personal computer installed in a patient's room was evaluated by Chan et al. (2002). This smart tool system proved valid in assessing and recording data on activities such as getting out of bed, mobility and travel patterns of a psychotic patient with moderate cognitive decline and behavioral disorders.

Global Positioning Systems (GPS) can also assist in locating people with dementia. Recently, Miskelly (2005) tested a mobile phone with GPS to help families of dementia sufferers if their relative happens to wander off. The person with dementia carries the mobile phone when leaving the house. If a carer or relative needs to know his or her whereabouts, a 24-h control centre can be called. By getting the coordinates of the phone they can pinpoint the person's location with an accuracy of 5 m, enabling help to be sent quickly.

The suitability of GPS systems for location tracking is highlighted by Shimizu et al. (2000) who developed a system that locates the wandering individual to a range of 100–200 m. The system was tested in various conditions such as adverse weather and close proximity of tall buildings. Results showed that despite these adverse conditions the location accuracy and frequency was sufficient for application in dementia wandering.

The Personal Handy-phone System, worn by the patient as a pendant, could transmit the patient's position with an accuracy of 60 m to a personal computer. The computer generates a map of the area, and then automatically sends this map via email to the carer who can view it on a mobile phone (Ogawa et al., 2004).

Applying a user-driven design approach Orpwood et al. (2005) developed and evaluated a smart home environment consisting of different services that aimed to assist people with dementia in various areas. During the night, for example, a lighting system was activated if a patient had left the bed. On returning to bed the lights would fade off after a few minutes. However, failure to return to bed after a pre-defined period of time would result in the communication system telling the patient that it is nighttime and they should go to bed. Furthermore, the system communicated to the patients when they approached an exit during the night and verbally reminded the patient to return to bed. Next to night monitoring, the system incorporated a cooking monitor service, detecting problems and potential dangers during the cooking process. During evaluations the cooker monitor worked well. However, it caused some irritation to users when the cooker was turned off under false positive situations.

The 'eHealth Strategic Objective' of the Information Society Technologies research program of the European Commission is aiming to create an 'intelligent environment' that allows ubiquitous management of each person's health status, assisting patients and health professionals in coping with major health challenges. The focus is on key technologies, such as biosensors and secure communications in 'smart clothes' and implants, as well as software tools for monitoring and managing health status and patient safety. As far as we know such technologies are not yet being applied and validated for persons with dementia.

#### 4. Conclusion and discussion

The aim of this review was to get an overview of ICT solutions that successfully support people with dementia in the areas of obtaining generalized and personal information, coping with symptoms of dementia, maintaining social contact and company, and enhancing (feelings of) safety.

##### 4.1. Available ICT-based services per needs area

Based on this literature study we can conclude that digital *information* on dementia, health companies, services and support offerings or financial and legal issues is provided with varying quality on multiple websites, as well as by digital social charts. A majority of these sites offer useful tips and support for carers but seem less attuned to the person suffering from dementia. This applies to the web design as well as to the information offered. Unfortunately, we did not find research in which the websites were tested on helpfulness for the target group. Additionally, an overriding drawback of all digital information is its generic nature. Demand-orientated, personalized information is still difficult to obtain. As the need for personalized and context-sensitive information clearly exists (Roest et al., 2007, submitted), future development of digital information systems should take these aspects into account.

In the last decade various ICT solutions were developed to support people with dementia and their carers in their everyday problems that arise from the *symptoms* associated with dementia. External memory aids in the form of reminders (to take medication or keep appointments) or cognitive interventions with Internet-based multimedia systems, including familiar equipment such as TV or PC, have proven efficacious in stimulating memory as well as enhancing feelings of independence and autonomy in the person with dementia. Several devices were developed to compensate for cognitive disabilities in the ENABLE project (e.g. a cooker usage monitor). All in all, these studies show that persons with dementia are not only capable of handling electronic equipment, but also benefit in terms of more confidence and enhanced positive affect, thereby indirectly reducing the carer's perceived burden. Information and emotional support for carers in coping with dementia symptoms furthermore is available through web-forums and chat-rooms, video-training and telephone services. Though web-based interventions seem promising especially to help meeting the various needs of people with dementia and carers in an effective and less time-consuming way, instrumental ICT-support for coping with behavioral and psychological changes in dementia is, however, relatively disregarded as yet. People with dementia and their carers have access to online information (e.g. site of the Alzheimer Association), support (e.g. 'Carer's Friend'; Beauchamp et al., 2005), education (e.g. AlzOnline's Positive Care giving classes; Glueckauf et al., 2004) and advice (e.g. online consultation of psychiatrist on <http://www.hulpgids.nl>) on coping with behavioral and psychological changes. However, apart from a study in which recorded music in combination with sung messages was found to be highly effective in guiding the behavior of people with dementia

(Yasuda et al., 2006), no proven successful instrumental ICT solutions for coping with behavioral and psychological changes were found. The integration of different technologies into a combined modular system, as piloted in several European projects such as FP4-TASC and FP5-TeleCARE, looks promising. Patients did not only made less use of conventional health services, but they also reported less negative feelings. Further effort into integrating different systems into a service bundle delivered to patients is therefore recommended.

Support for *social contact and company* is realized through simplified mobile phones or videophones, that have been reported to facilitate communication between people with dementia and their family or friends. Enhanced positive affect as well as increased activity and communication levels have been observed with computer software providing music or video memories or robotics, such as a toy dog or an entertainment robot. These technical solutions proved to be easily implemented and of great benefit to the person with dementia and their carers.

Finally, we found several ICT solutions for *health monitoring and to enhance perceived safety*. GPS technology for tracking wandering or lost persons as well as monitoring systems to detect fire or gas leakage or signal nighttime activity allow for unobtrusive yet efficient assessments of safety. Additionally, people with dementia and their carers report enhanced feelings of safety and security and less fear and anxiety. Dissemination and implementation of these systems into households is therefore advisable. The monitoring of biological functions in people with dementia is still in its infancy. Unobtrusive, ubiquitous monitoring and management of a person's health using an 'intelligent environment' (e.g. the 'eHealth Strategic Objective') seems to have potential for the future but has yet to prove its efficacy in the field.

#### 4.2. Limitations of the current study

In order to make the review study practical a number of limitations are present both in the searching process and the analysis. A first restriction concerned the focus of the study. Only ICT solutions that could contribute to meet the most frequently mentioned *unmet* needs by people with dementia and their informal carers (see Table 1) were included in the study. ICT-solutions in other needs areas were therefore not included in this study.

A second restriction limited the searching process to a number of databases which covered both the clinical and technical aspects of the study. The databases PubMed, PsychInfo, Cinahl, INSPEC, IEEE and Google (Scholar) were used in the searching process with a total of 165 articles found. Although this selection may have caused us to miss relevant articles, we feel the variety of articles found provides a good overview of the state of the art of ICT solutions for the selected needs of people with dementia.

A third restriction was that we included only articles that reported on ICT solutions that were proven to be helpful for people with dementia and/or their carers and that were published before 15 November 2006. ICT solutions that are currently on the market and potentially useful for the target group, but for which no publications were found on the usefulness for people with dementia and/or their carers, are therefore not included in this review.

#### 4.3. Societal relevance

ICT-support for needs that are unmet by the present care and welfare services could optimize the care for people with dementia and their carers and thus enhance their quality of life.

Whether people with dementia or their informal carers actually make use of and benefit from available ICT-applications and services will partly depend on the attitude of health care professionals towards these solutions (e.g. Larner, 2003). People with dementia and their informal carers are most likely to learn about available ICT solutions through their General Practitioner, specialist, insurance company or care and welfare organizations. In order to inform their clients well, health care professionals should be aware of current and new ICT developments and services. This review gives an overview of successfully and partially successfully applied ICT solutions for people with dementia and their carers and therefore can help health care professionals to get informed on this subject and transfer this knowledge to their clients.

Not all ICT-applications and services on today's market have been tested in advance or applied successfully for people with dementia. Although some untested systems could be beneficial to people with dementia or their carers, one cannot be sure of the efficacy of these systems. Only when the system is installed and in use, will the actual effect become clear, in the worst case causing financial, administrative or emotional difficulties for the end-user. Without pretending to be complete, this review tried to focus specifically on ICT solutions that have been successfully tested on

people with dementia or their informal carers and have proven to be helpful. This review can therefore support professionals, people with dementia and their informal carers in selecting a specific solution. The information gathered in this review can also be used by health care professionals, policy makers and account managers of insurance companies to assess whether an application or service should be included in National Health Service packages or should be subsidized from healthcare funds.

As people with mild to moderate dementia seem capable of expressing their needs and do so from a different perspective than their informal carer (Dröes et al., 2006; Roest et al., 2007, submitted), ICT solutions should be developed in dialogue with the target-users. Feedback from people with dementia and their carers during the development process can be used to make the final product better attuned to their needs, more personalized and more helpful (COGKNOW Consortium, 2006; Sixsmith et al., 2007).

#### 4.4. Technological interest and research recommendations

Previous research on ICT solutions that aim to meet the unmet needs of people with dementia and their carers in the need areas we focused on in this review study primarily focused on two areas. Most studies concerned general information on the disease, how to cope with the consequences, and service offerings on the one hand, and on the safety and monitoring of the person with dementia in various situations on the other hand. The majority of the latter solutions were particularly concerned with stray prevention and adopted a number of varying techniques to detect or locate the person with dementia when wandering both within and outside of the home environment. In the majority of solutions for stray prevention and detection, a warning system was adopted to alert carers or close family members to particular situations that could pose a risk to the person with dementia. Another area that the research explored was that of social contact and providing a basis for allowing a person with dementia to communicate better with others. The research articles found in this area highlighted the need for technology to allow persons with dementia to improve on their communication skills with others. Two studies showed that by using videophone technology, the quantity and quality of conversations between the person with dementia and others would be enhanced (Mickus and Luz, 2002; Savenstedt et al., 2003).

One area that was relatively poorly represented was support for persons with dementia to compensate for their disabilities. Although quite a number of solutions were found for memory problems that could potentially be applicable to person with dementia, for some there was no direct evidence that they were effective in the target group. Therefore, these systems were excluded from the review. In a similar fashion there were a number of systems that aimed to provide help with daily living activities such as brushing your teeth; however, these solutions were never tested on people with dementia and were therefore also excluded from the review. Though successfully applied in a single study (Yasuda et al., 2006), instrumental ICT-support for coping with behavioral and psychological symptoms in dementia is relatively disregarded as yet and the same holds for personalized information on the diagnosis, condition and personal care appointments. In summary it can be noted that there is a lack of ICT research conducted for two of the four areas that this review process addresses, namely support for obtaining personalized information and instrumental support for dealing with symptoms of dementia, such as memory problems, behavioral and psychological changes, and problems with daily activities.

It is recommended that future studies should concentrate on integrating and applying current techniques and solutions in real life situations with persons with dementia. This research and development process should be user-driven “to ensure that devices and systems are grounded within a thorough understanding of the needs, preferences and desires of potential users.” (Sixsmith et al., 2007). Special attention should be given to ICT solutions that provide personalized information; compensate for disabilities such as memory problems; help people with dementia and their carers to cope with the behavioral and psychological changes associated with the dementia; and provide help with daily activities, including pleasant activities (Dröes et al., 2006; Sixsmith et al., 2007). The former two need particular attention in producing new ideas on how technology can be of assistance for the target group. An example of a personalized information service is the dynamic interactive social chart for dementia care (DEM-DISC) that provides people with dementia and their informal and formal carers with tailor-made information on regional and national care and support services (and service bundles) (Dröes et al., 2005a,b). ICT solutions for helping people to cope with behavioral and psychological changes and help with daily activities require selection of existing smart home solutions and the subsequent adaptation and testing of them for persons with dementia. All of the authors of this review participate or are indirectly involved in the COGKNOW project (2006) that aims at helping people with mild dementia

navigate through their day. The project focuses on developing integrated ICT solutions for frequently mentioned unmet needs by people with dementia and their carers (Dröes et al., 2006; Roest et al., 2007, submitted), including memory support, support for social contact and daily activities, and support for enhancing feelings of safety.

The majority of the studies found included small user groups and were uncontrolled, which makes it hard to generalize their outcomes. Therefore, a second recommendation would be to test the effectiveness of ICT solutions on a larger scale in, preferably randomized controlled trials before implementing them in the care for people with dementia.

## Acknowledgements

The study was conducted within the framework of two projects: the FReeband User eXperience (FRUX) project that was financed by the Dutch Ministry of Economic Affairs under contract BSIK 03025, Dioraphte Foundation, RCOAK, NHDI, Foundation het Zonnehuis and Stichting Alzheimer & Neuropsychiatrie Foundation and the COGKNOW project that is funded by the European Commission's Information Society Technologies (IST) program under grant 034025.

## References

- Ager, A., Aalykke, S., 2001. TASC: A microcomputer support system for persons with cognitive disabilities. *Br. J. Educat. Technol.* 32, 373–377.
- Algase, D.L., Beattie, E.R.A., Leitsch, S.A., Beel-Bates, C.A., 2003. Biomechanical activity devices to index wandering behaviour in dementia. *Am. J. Alzheimer's Dis. Other Demen.* 18, 85–92.
- Alm, N., Dye, R., Gowans, G., Campbell, J., Astell, A., Ellis, M., 2003. Designing an interface usable by people with dementia. *ACM SIGCAPH Computers and the Physically Handicapped*, June–September, pp. 156–157.
- Baruch, J., Downs, M., Baldwin, C., Bruce, E., 2004. A case study in the use of technology to reassure and support a person with dementia. In: Moriarty, J. (Ed.), *Dementia. Innovative Practice*, vol. 3 (3). SAGE Publications, USA, pp. 371–392.
- Beauchamp, N., Irvine, A.B., Seeley, J., Johnson, B., 2005. Worksite-based Internet multimedia program for family caregivers of persons with dementia. *Gerontologist* 45, 793–801.
- Biegel, D.E., Bass, D.M., Schulz, R., Morycz, R., 1993. Predictors of in-home and out-of-home service use by family caregivers of Alzheimer's disease patients. *J. Aging Health* 5, 419–438.
- Blackburn, P., 1988. Freedom to wander. *Nurs. Times* 84, 54–55.
- Bourgeois, M., Dijkstra, K., Burgio, L., Allen-Burge, R., 2001. Memory aids as an augmentative and alternative communication strategy for nursing home residents with dementia. *Augmentative Altern. Commun.* 17, 196–210.
- Brodsky, H., Thomson, C., Thompson, C., Fine, M., 2005. Why caregivers of people with dementia and memory loss do not use services. *Int. J. Geriatr. Psychiatry* 20, 537–546.
- Broek, M.D.v.d., Downes, J., Johnson, Z., Dayus, B., Hilton, N., 2000. Evaluation of an electronic memory aid in the neuropsychological rehabilitation of prospective memory deficits. *Brain Injury* 14, 455–462.
- Burns, A., 2000. The burden of Alzheimer's disease. *Int. J. Neuropsychopharmacol.* 3, 31–38.
- Camarinha-Matos, L., Asfaramesh, H., 2004. TeleCARE: Collaborative virtual elderly care support communities. *J. Inf. Technol. Healthc.* 2, 73–86.
- Chan, M., Campo, E., Laval, E., Estève, D., 2002. Validation of a remote monitoring system for the elderly: Application to mobility measurements. *Technol. Health Care* 10, 391–399.
- Clare, L., Wilson, B.A., Carter, G., Breen, K., Gosses, A., Hodges, J.R., 2000. Intervening with everyday memory problems in dementia of Alzheimer type: an errorless learning approach. *J. Clin. Exp. Neuropsychol.* 22, 132–146.
- Clare, L., 2002. We'll fight it as long as we can: coping with the onset of Alzheimer's disease. *Aging Ment. Health* 6, 139–148.
- Coe, M., Neufeld, A., 1999. Male caregivers' use of formal support. *West. J. Nurs. Res.* 21, 568–588.
- COGKNOW Consortium, 2006. COGKNOW; helping people with dementia navigate their day. Description of Work, Project Proposal, EU Contract No. 034025.
- Cox, C., 1997. Findings from a state-wide program of respite care: a comparison of service users, stoppers, and nonusers. *Gerontologist* 37, 511–517.
- Cummings, J.L., Mega, M., Gray, K., Rosenberg-Thompson, S., Carusi, D.A., Gornbein, J., 1994. The Neuropsychiatric Inventory: comprehensive assessment of psychopathology in dementia. *Neurology* 44, 2308–2314.
- Dröes, R.M., Lindeman, E.M., Breebaart, E., Tilburg, W.v., 1996. Determinanten van belasting van verzorgers van mensen die lijden aan dementie. In: Dröes, R.M. (Ed.), *Amsterdams Ontmoetingscentra, een nieuwe vorm van ondersteuning voor mensen met dementie en hun verzorgers. Eindrapport 1996*. Thesis Publishers, Amsterdam, pp. 89–118.
- Dröes, R.M., Hout, H.P.I.van., Ploeg, E.S.van.de., 2004. Camberwell Assessment of Need for the Elderly (CANE). Revised Version IV. Nederlandse vertaling. VU medisch centrum, Amsterdam.
- Dröes, R.M., Meiland, F.J.M., Doruff, C., Varodi, I., Akkermans, H., Baida, Z., Faber, E., Haaker, T., Moelaert, F., Kartseva, V., Tan, Y.H., 2005a. A dynamic interactive social chart in dementia care. Attuning demand and supply in the care for persons with dementia and their carers. In: Bos, L., Laxminarayan, S., Marsh, A. (Eds.), *Medical and Care Compunctics, Studies in Health Technology and Informatics 2*[114]. IOS Press, The Netherlands, USA, England, pp. 210–220.

- Dröes, R.M., Meiland, F.J.M., Roest, H.G.v.d., Marocchini, R., Slagter, R.S., Baida, Z., Haaker, T., Kartseva, V., Hulstijn, J., Schmieman, R., Akkermans, H., Faber, E., Tan, Y.H., 2005b. Opportunities for we-centric service bundling in dementia care. Project Freeband/FRUX health care pilot report D1.7.1-II, D1.7.2-II, D1.7.3-II, D2.5.1-II, D2.5.2-II, D2.5.4-II, Amsterdam.
- Dröes, R.M., Boelens, E.J., Bos, J., Meihuizen, L., Ettema, T.P., Gerritsen, D.L., Hoogeveen, F., Lange, J.de., Schölzel-Dorenbos, C., 2006. Quality of life in dementia in perspective; an explorative study of variations in opinions among people with dementia and their professional caregivers, and in literature. *Demen. Int. J. Soc. Res. Pract.* 5, 533–558.
- Duijnste, M., 1992. De belasting van familieleden van dementerenden. Intro, Nijkerk.
- Dunkin, J.J., Anderson-Hanley, C., 1998. Dementia caregiver burden: a review of the literature and guidelines for assessment and intervention. *Neurology* 51, S53–S60.
- Ferri, C.P., Prince, M., Brayne, C., Brodaty, H., Fratiglioni, L., Ganguli, M., Hall, K., Hasegawa, K., Hendrie, H., Huang, Y., Jorm, A., Mathers, C., Menezes, P.R., Rimmer, E., Sczufca, M., 2005. Global prevalence of dementia: a Delphi consensus study. *Lancet* 366, 2112–2117.
- Freeman, E., Clare, L., Savitch, N., Royan, L., Litherland, R., Lindsay, M., 2005. Improving website accessibility for people with early stage dementia: a preliminary investigation. *Aging Ment. Health* 9, 442–448.
- Fritschy, E.P., Kessels, R.P., Postma, A., 2004. External memory aids for patients with dementia: a literature study on efficacy and applicability. *Tijdschr. Gerontol. Geriatr.* 35, 234–239.
- Gilliard, J., Hagen, I., 2004. Enabling technologies for people with dementia. Cross-national analysis report. D4.4.1. QLK6-CT-2000-00653, pp. 1–69.
- Glueckauf, R.L., Ketterson, T.U., Loomis, J.S., Dages, P., 2004. Online support and education for dementia caregivers: Overview, utilization, and initial program evaluation. *Telemed. J. e-Health* 10, 223–232.
- Grandmaison, E., Simard, M., 2003. A critical review of memory stimulation programs in Alzheimer's disease. *J. Neuropsychiatry Clin. Neurosci.* 15, 130–144.
- Gwyther, L.P., 1998. Social issues of the Alzheimer's patient and family. *Am. J. Med.* 104, 17S–21S.
- Hancock, G.A., Woods, B., Challis, D., Orrell, M., 2006. The needs of older people with dementia in residential care. *Int. J. Geriatr. Psychiatry* 21, 43–49.
- Hart, T., Buchhofer, R., Vaccaro, M., 2004. Portable electronic devices as memory and organizational aids after traumatic brain injury: a consumer survey study. *J. Head Trauma Rehabil.* 19, 351–365.
- Hassol, A., Walker, J.M., Kidder, D., Rokita, K., Young, D., Pierdon, S., Deitz, D., Kuck, S., Ortiz, E., 2004. Patient experiences and attitudes about access to a patient electronic health care record and linked web messaging. *J. Am. Med. Inform. Assoc.* 11, 505–513.
- Health Council of the Netherlands, 2002/04. Dementia. The Hague, Health Council of the Netherlands.
- Hersh, N.A., Treadgold, L.G., 1994. NeuroPage: the rehabilitation of memory dysfunction by prosthetic memory and cueing. *Neurorehabilitation* 4, 187–197.
- Hofmann, M., Hock, C., Kuhler, A., Muller-Spahn, F., 1996. Interactive computer-based cognitive training in patients with Alzheimer's disease. *J. Psychiatry Res.* 30, 493–501.
- Holte, T., Hagen, I., Björneby, S., 1998. Evaluation of an electronic calendar as helping aid for persons suffering from memory problems or cognitive impairment. Report of the TED-group.
- Honeyman, A., Cox, B., Fisher, B., 2005. Potential impacts of patient access to their electronic care records. *Inform. Prim. Care* 13, 55–60.
- Ikarashi, A., Magara, K., Nonaka, S., Ohno, H., 2002. The searching system for wandering demented aged person using GPS. *Trans. Inst. Electr. Eng. Jpn.* 122, 609–616.
- Inglis, E.A., Szymkowiak, A., Gregor, P., Newell, A.F., Hine, N., Shah, P., Wilson, B.A., Evans, J., 2003. Issues surrounding the user-centred development of a new interactive memory aid. *Universal Access Inf. Soc.* 2, 226–234.
- Kim, H.J., Burke, D.T., Dowds Jr., M.M., Boone, K.A., Park, G.J., 2000. Electronic memory aids for outpatient brain injury: follow-up findings. *Brain Injury* 14, 187–196.
- Kinney, J., Kart, C.S., Murdoch, L., Conley, C., 2004. Striving to provide safety assistance for families of elders: the SAFE House project. *Demen. Int. J. Soc. Res. Pract.* 3, 351–370.
- Koffman, J., Taylor, S., 1997. The needs of caregivers. *Elder Care* 9, 16–19.
- Kort, S., 2005. Mobile Coaching. A pilot study into the user-friendliness and effects of Mobile Coaching on the wellbeing of people with dementia and their informal caregivers. Faculty of Psychology, Vrije Universiteit, Amsterdam.
- Larner, A.J., 2003. Use of the Internet and of the NHS direct telephone helpline for medical information by a cognitive function clinic population. *Int. J. Geriatr. Psychiatry* 18, 118–122.
- Lekeu, F., Wojtasik, V., Linden, M.v.d., Salmon, E., 2002. Training early Alzheimer patients to use a mobile phone. *Acta Neurol. Belg.* 102, 114–121.
- Lin, C.C., Chiu, M.J., Hsiao, C.C., Lee, R.G., Tsai, Y.S., 2006. Wireless health care service system for elderly with dementia. *IEEE Trans. Inf. Technol. Biomed.* 10, 696–704.
- Lindström, J.I., 2005. Navigation, alarming and positioning. A preliminary study conducted in Sweden by the Royal Institute of Technology (KTH). Department of Speech, Music and Hearing, on the assignment of the National Post and Telecom agency (PTS) (PTS-ER-2006-16).
- Long, C.O., Williams, J.K., 2005. Exploring Alzheimer's disease on the Web. *Nursing* 35, 30–31.
- Magnusson, L., Berthold, H., Chambers, M., Brito, L., Emery, D., Daly, T., 1998a. Using telematics with older people: the ACTION project. Assisting carers using telematics interventions to meet older persons' needs. *Nurs. Stand.* 13, 36–40.
- Magnusson, L., Berthold, H., Brito, L., Chambers, M., Emery, D., Daly, T., 1998b. ACTION: Assisting Carers using Telematics Interventions to Meet Older Persons' Needs. 28-8-0098.
- Magnusson, L., Hanson, E., Brito, L., Berthold, H., Chambers, M., Daly, T., 2002. Supporting family carers through the use of information and communication technology—the EU project ACTION. *Int. J. Nurs. Stud.* 39, 369–381.

- Masuda, Y., Yoshimura, T., Nakajima, K., Nambu, M., Hayakawa, T., Tamura, T., 2002. Unconstrained monitoring of prevention of wandering the elderly. In: *Engineering in Medicine and Biology, 24th Annual Conference and the Annual Fall Meeting of the Biomedical Engineering Society (EMBS/BMES Conference)*. Proceedings of the Second Joint 2. pp. 1906–1907.
- Mickus, M.A., Luz, C.C., 2002. Televisits: sustaining long distance family relationships among institutionalized elders through technology. *Aging Ment. Health* 6, 387–396.
- Miskelly, F., 2004. A novel system of electronic tagging in patients with dementia and wandering. *Age Ageing* 33, 304–306.
- Miskelly, F., 2005. Electronic tracking of patients with dementia and wandering using mobile phone technology. *Age Ageing* 34, 497–499.
- Ogawa, H., Yonezawa, Y., Maki, H., Sato, H., Caldwell, W.M., 2004. A mobile phone-based safety support system for wandering elderly persons. In: *26th Annual International Conference of the Engineering in Medicine and Biology Society (EMBC)*. Conference Proceedings 2. pp. 3316–3317.
- Olsen, R.V., Hutchings, B.L., Ehrenkrantz, E., 2000. Media Memory Lane™ interventions in an Alzheimer's day care centre. *Am. J. Alzheimer's Dis. Other Demen.* 15, 163–175.
- Oriani, M., Moniz-Cook, E., Binetti, G., Zanieri, G., Frisoni, G.B., Geroldi, C., Vreese, L.P.d., Zanetti, O., 2003. An electronic memory aid to support prospective memory in patients in the early stages of Alzheimer's disease: a pilot study. *Aging Ment. Health* 7, 22–27.
- Orpwood, R., Gibbs, C., Adlam, T., Faulkner, R., Meegahawatte, D., 2005. The design of smart homes for people with dementia—user-interface aspects. *Universal Access Inf. Soc.* 4, 156–164.
- Paavilainen, P., Korhonen, I., Lotjonen, J., Cluitmans, L., Jylha, M., Sarela, A., Partinen, M., 2005. Circadian activity rhythm in demented and non-demented nursing-home residents measured by telemetric actigraphy. *J. Sleep. Res.* 14, 61–68.
- Perry, J., Bontinen, K., 2001. Evaluation of a weekend respite program for persons with Alzheimer disease. *Can. J. Nurs. Res.* 33, 81–95.
- Poon, P., Hui, E., Dai, D., Kwok, T., Woo, J., 2005. Cognitive intervention for community-dwelling older persons with memory problems: telemedicine versus face-to-face treatment. *Int. J. Geriatr. Psychiatry* 20, 285–286.
- Proctor, R., Testad, I., 2005. Carer stress—overview. In: Burns, A. (Ed.), *Standards in Dementia Care: European Dementia Consensus Network* Edcon. Taylor & Francis Group, New York, pp. 241–273.
- Roest, H.G.v.d., Meiland, F.J.M., Marocchini, R., Comijs, H.C., Jonker, C., Dröes, R.M., 2007. Subjective needs of people with dementia: a review of the literature. *Int. Psychogeriatr.* 19 (June (3)), 559–592.
- Roest, H.G.v.d., Meiland, F.J.M., Comijs, H.C., Jansen, A.P., Hout, H.P.J.van. What do you need? That is the question. The subjective and objective needs of people with dementia living in the community, submitted.
- Sakairi, K., 2004. Research of robot-assisted activity for the elderly with senile dementia in a group home. In: *SICE 2004 Annual Conference* 3. pp. 2092–2094.
- Savenstedt, S., Brulin, C., Sandman, P.O., 2003. Family members' narrated experiences of communicating via video-phone with patients with dementia staying at a nursing home. *J. Telemed. Telecare* 9, 216–220.
- Shimizu, K., Kawamura, K., Yamamoto, K., 2000. Location system for dementia wandering. In: *22nd Annual International Conference of the IEEE Engineering in Medicine and Biology Society*. Proceedings of the 22nd Annual International Conference of the IEEE 2. pp. 1556–1559.
- Sixsmith, A.J., Gibson, R.D., Orpwood, R.D., Torrington, J.M., 2007. Developing a technology 'wish-list' to enhance the quality of life of people with dementia. *Gerontechnology* 6, 2–19.
- Szymkowiak, A., Morrison, K., Shah, P., Gregor, P., Evans, J., Newell, A.F., Wilson, B.A., Shofield, S., 2004. Memojog—an interactive memory aid with remote communication. In: Keates, S., Clarkson, J., Langdon, P., Robinson, P. (Eds.), *Designing a more inclusive world*. Proceedings of Second Cambridge Workshop on Universal Access and Assistive Technology (CWUAAT), 22 April, Cambridge, UK, pp. 15–24.
- Tamura, T., Yonemitsu, S., Itoh, A., Oikawa, D., Kawakami, A., Higashi, Y., Fujimooto, T., Nakajima, K., 2004. Is an entertainment robot useful in the care of elderly people with severe dementia? *J. Gerontol. A Biol. Sci. Med. Sci.* 59, 83–85.
- Tarraga, L., Boada, M., Modinos, G., Espinosa, A., Diego, S., Morera, A., Guitart, M., Balcells, J., Lopez, O.L., Becker, J.T., 2006. A randomised pilot study to assess the efficacy of an interactive, multimedia tool of cognitive stimulation in Alzheimer's disease. *J. Neurol. Neurosurg. Psychiatry* 77, 1116–1121.
- Vehvilainen, L.M., Zielstorff, R., Gertman, P.M., Tzeng, M.C., Estey, G., 2002. Alzheimer's caregiver Internet support system (ACISS): evaluating the Feasibility and effectiveness of supporting family caregivers virtually. In: *American Medical Informatics Association 2002 Symposium*. 11 November (S32 Poster Session 1).
- White, M.H., Dorman, S.M., 2000. Online support for caregivers. *Analysis of an Internet Alzheimer mailgroup*. *Comput. Nurs.* 18, 168–176.
- Wilson, B.A., Emslie, H.C., Quirk, K., Evans, J.J., 2001. Reducing everyday memory and planning problems by means of a paging system: a randomised control crossover study. *J. Neurol. Neurosurg. Psychiatry* 70, 477–482.
- Winslow, B.W., 2003. Family caregivers' experiences with community services: a qualitative analysis. *Public Health Nurs.* 20, 341–348.
- Woolham, J., 2005. Safe at Home. In: *The effectiveness of assistive technology in supporting the independence of people with dementia: the Safe at Home project*, Hawker Publications, London.
- Yasuda, K., Beckman, B., Yoneda, M., Yoneda, H., Iwamoto, A., Nakamura, T., 2006. Successful guidance by automatic output of music and verbal messages for daily behavioural disturbances of three individuals with dementia. *Neuropsychol. Rehabil.* 16, 66–82.
- Zanetti, O., Zanieri, G., Vreese, L.P.d., Frisoni, G.B., Binetti, G., Trabucchi, M., 2000. Utilizing an electronic memory aid with Alzheimer's disease patients. A study of feasibility. In: Paper presented at the Sixth International Stockholm/Springfield Symposium on Advances in Alzheimer Therapy.